

What's on the menu? A presumed attack of Andean bear on a Mountain tapir at the Puracé National Natural Park, Colombia

Gustavo Adolfo Pisso-Florez^{1,2}, Ignacio Gómez-Lora³, I. Mauricio Vela-Vargas^{4,5}, Héctor Pizo¹, Isaac Bedoya Dorado¹, Héctor E. Ramírez-Chaves⁶

1 Parque Nacional Natural Puracé, Carrera 9 # 25 N – 06, Popayán, Cauca, Colombia

2 Fundación Biotellus, Carrera 11 # 5-43, Popayán, Cauca, Colombia

3 Fundación Zoológica de Cali, Carrera. 2 west, Calle 14, Cali, Colombia

4 Wildlife Conservation Society – Colombia, Carrera. 13 No. 77 a- 42, Bogotá, Colombia

5 Proyecto de Conservación de Aguas y Tierras – ProCAT Colombia. Carrera 11 # 96-43, Bogotá, Colombia

6 Departamento de Ciencias Biológicas, Facultad de Ciencias Exactas y Naturales, Universidad de Caldas, and Centro de Museos, Museo de Historia Natural, Universidad de Caldas. Calle 65 # 26-10, Manizales, Caldas

Corresponding author: Gustavo Adolfo Pisso-Florez (tapiflo@gmail.com)

Academic editor: A.M. Leal-Zanchet | Received 6 August 2020 | Accepted 19 December 2020 | Published 14 January 2021

Citation: Pisso-Florez GA, Gómez-Lora I, Vela-Vargas IM, Pizo H, Bedoya Dorado I, Ramírez-Chaves HE (2021) What's on the menu? A presumed attack of Andean bear on a Mountain tapir at the Puracé National Natural Park, Colombia. *Neotropical Biology and Conservation* 16(1): 19–25. <https://doi.org/10.3897/neotropical.16.e57140>

Abstract

Two iconic and charismatic species that inhabit the northern Andes of South America are the Andean bear (*Tremarctos ornatus*) and the Mountain tapir (*Tapirus pinchaque*). Both species can be found sympatrically in several areas of Colombia, Ecuador, and northern Peru. Despite their overlap in distribution, little is known about interactions between both species, with few reported cases of Andean bear attacks on the Mountain tapir. Here, we report a possible attack by an Andean bear on a Mountain tapir in the northern part of Puracé National Natural Park, Colombia based on strong wounds and marks on a tapir's back and rump. The wounds match typical attack patterns generated by Andean bears and corroborates previous camera traps records of bears attacking tapirs in this locality.

Keywords

ecological interactions, neotropical mammals, high Andean forest, protected area, *Tapirus pinchaque*, *Tremarctos ornatus*

The Mountain tapir *Tapirus pinchaque* (Roulin 1829) is the smallest extant species of the genus *Tapirus* and has restricted distribution to cloud forest remnants and páramo habitats in Colombia, Ecuador and northeastern Peru, at elevations between 1400 and 4700 m (Lizcano et al. 2002; Arias-Alzate et al. 2010; Cavelier et al. 2010; Padilla et al. 2010; Ortega-Andrade et al. 2015). Mountain tapirs are among the largest mammals of the north Andean region reaching a weight up to 190 kg (Lizcano et al. 2017). Within its range, the Mountain tapir is sympatric with the Andean bear, *Tremarctos ornatus* Cuvier, 1825, described as one of its potential predators (Castellanos 2011, 2019; Castellanos 2014; Rodríguez et al. 2014). Andean bears are considered as an omnivorous species with a high preference for plants, and animal proteins comprise less than 10% of its diet (Figueroa 2013; Gonzales et al. 2016). Evidence of predatory attempts has been reported based on wildlife camera photographs observing an Andean bear holding on to the back to a Mountain tapir (Rodríguez et al. 2014). Besides scavenging records, the presence of Mountain tapir hair in Andean bear feces and claw marks on mountain tapir individuals have been suggested as further evidence of interactions between these two species (Castellanos 2011; Rodríguez et al. 2014; Castellanos 2019; López-Ordóñez et al. 2020). Here, we present a report of an individual Mountain tapir with wound patterns typically generated by Andean bear attack at the Puracé National Natural Park (PNNP), southwestern Colombia.

On 21 July 2020, local people reported to rangers of the Puracé National Natural Park (PNNP) the presence of wounded Mountain tapir around the Güargüero stream in the sector Alto Vedón La Plata, 39–40 km of the road Puracé – La Plata (02°19'33.8"N, 076°17'19.6"W \pm 25–30 m; 2,890 m a.s.l.) at the PNNP, Department of Cauca, southwestern Colombia (Fig. 1). This area is mainly covered by High-Andean cloud forests and páramo ecosystems, in the southern part of the Central Cordillera of Colombia (Bonilla-Valencia et al. 2019). Three PNNP rangers tracked the tapir and made observations of the individual for five days (23 to 27 July 2020) along seven km of road in the protected area. During 23 July, the survey was made between 14h:30–17h:00. Similarly, on 24 July, two surveys were made between 9h:00–11h:00 and 15h:00–17h:00, and on 25–27 July, three surveys were made between 15h:00 and 19h:00, all on the locality reported by local communities. All surveys were diurnal and occurred in the rainy season (Heck and Figueroa 2009), however no precipitation was recorded during the surveys. Our efforts included a 55.5 hours/person and a total of 35 km of road sampled.

On 24 July 2020, we first observed the wounded Mountain tapir, a juvenile individual (Fig. 2A) with two large wounds located in the right flank on the anterior part of the back ($\sim 22 \times 16$ cm), a wound located on the posterior region




Figure 1. Locality of wounded Mountain tapir record (yellow star) at the Puracé National Natural Park, Colombia.

($\sim 11 \times 5$ cm; Fig. 2B) and on the right side of its rump ($\sim 17 \times 12$ cm; Fig. 2C), and small scratches on the rump and both thighs (Fig. 2D), which we presume were made by the Andean bear.

The same day at 16h:00, one of the rangers observed the wounded Mountain tapir feeding along the road edge for one hour and was able to collect a rotten piece of meat hanging from the wound of the rump, after stored in 70% ethanol (personal field number – GAP 198). The tapir exhibited abnormal movements with its right front foot but remained upright. The individual eventually entered the forest.

On 25 July at 17h:48, the PNNP team observed the individual 600 m away from its last sighting, foraging on the roadside for approximately 20 minutes. The tapir exhibited no signs of discomfort, such as rubbing its head or scratching branches into the wounds or difficulty walking. The individual again entered the forest around 18h:10. Two days later on 27 July, the PNNP team detected the individual 1.4 kilometers away from the last observed location. The dry wounds were about 60% of the size since the first observation. We attempted to record more precise measurements, but the arriving of local bus spooked the animal that into the surrounding forest. No capture or veterinary intervention to the animal were performed because it was



Figure 2. Records of a wounded Mountain tapir produced presumably by Andean bear at the Puracé National Natural Park, Colombia on 27 July 2020. **A** General view of the individual, **B** wound located in the right flank on the anterior part of the back, **C** wound located on the right rump region, **D** scratch located on the left thigh region. Photographs: Gustavo Adolfo Pisso-Florez.

standing and feeding normally. Also, the contingency measures taken by Colombian government because COVID-19 pandemic limited the mobilization of technical personnel to the PNNP. Also, the tapir was feeding normally and showed signs of healing naturally.

We suggest that the wounds observed on the Mountain tapir were caused by an Andean bear based on the patterns registered of bears attacking livestock (Goldstein et al. 2006; Figueroa 2015). Andean bears attacks large prey, by grasping them by the back with their claws and biting the back to disable it. This process produces multiple bites that leave large wounds and tears of flesh on the shoulders, back, and chest (Goldstein et al. 2006; Castellanos 2011; Narvaez and Zapata-Ríos 2016). Recent evidence has shown that the Andean bear preys on Mountain tapirs by hunting them (Castellanos 2019), or through scavenging (López-Ordóñez et al. 2020). The Mountain tapir wounds that we observed allow us to hypothesize that an Andean bear tried to attack the back of the tapir but failed to achieve the predation attempt (Rodríguez et al. 2014). Mountain tapirs are large, yet overtly cautious, and can move fast; such traits may allow them to escape attacking predators (Castellanos 2011; Rodríguez et al. 2014).

The road side behaviors of this Mountain tapir after a presumable Andean bear attack are similar to an event described in Ecuador (Castellanos 2019), where a Mountain tapir spent most of its time at nearby cattle pastures and salt licks, and periodically finding shade and shelter in patches of cloud forest. This individual may

have used an open roadway to better detect predators; it has been suggested that Andean bears may be affected by roads (Hernani-Lineros et al. 2020) but more evidence is needed. Another large predator inhabiting Puracé is the cougar, *Puma concolor* (Linnaeus, 1771), but this species primarily feeds on prey such Brocket deer (*Mazama rufina*) and Northern Pudu (*Pudu mephistophiles*) (Hernández-Guzmán et al. 2011). Further, cougars often attack the throat or neck of prey (Márquez and Goldstein 2014; Narváez and Zapata-Ríos 2016). More foraging and predator-prey dynamics information is warranted to better understand the ecology of charismatic northern Andean megafauna.

Acknowledgements

Gustavo Pisso, Hector Pizo and Isaac Bedoya thank the Parque Nacional Natural Puracé, Parques Nacionales Naturales de Colombia, and the Puracé indigenous community for their support during the field surveys. We thank the inter-municipal transporters by alerting us about the wounded tapir. Gustavo Adolfo Pisso-Florez and Héctor E. Ramírez-Chaves thank Rufford Small Grants (Grants 28471-2 and 29491-2) for support. All authors acknowledge Dr. Brian Blais from the University of Arizona for his comments on how to improve the quality of the manuscript.

References

- Arias-Alzate A, Downer C, Delgado VCA, Sánchez-Londoño J (2010) Un registro de tapir de montaña (*Tapirus pinchaque*) en el norte de la cordillera Occidental de Colombia. *Mastozoología Neotropical* 17(1): 111–116. <http://www.redalyc.org/articulo.oa?id=45713277011>
- Bonilla Valencia J, Roncancio-Duque N, Pisso-Florez GA (2019) Portafolio de proyectos de investigación del Parque Nacional Natural Puracé. Documento adjunto al Plan de manejo 2020–2024 del PNN Puracé. Dirección Territorial Andes Occidentales, Parques Nacionales Naturales de Colombia, Popayán-Cauca, 58 pp.
- Castellanos A (2011) Do Andean bears attack mountain tapirs? *International Bear News* 20: 41–42.
- Castellanos A (2014) Co-occurrence of Andean bear and Mountain tapir in the Papallacta region, Cayambe Coca National Park, Ecuador: A brief description. *International Bear News* 23(2): 20–21.
- Castellanos A (2019) The threat of road-kill to Andean tapirs: The case of ‘Jorgito’, the Andean tapir that lived beside the Quito-Amazon highway, Ecuador. *Boletín técnico* 14. Serie Zoológica 14–15: 1–5.
- Cavelier J, Lizcano D, Yerena E, Downer C (2010) The Mountain tapir (*Tapirus pinchaque*) and Andean bear (*Tremarctos ornatus*): Two charismatic, large mammals in South American tropical montane cloud forest. In: Scatena LA, Hamilton, LS (Eds) *Tropical Montane Forest: Science for Conservation and Administration*. Cambridge University Press, Cambridge, 172–189. <https://doi.org/10.1017/CBO9780511778384.019>

- Figuerola J (2013) Revisión de la dieta del Oso Andino *Tremarctos ornatus* (Carnivora: Ursidae) en América del Sur y nuevos registros para Perú. *Revista del Museo Argentino de Ciencias Naturales* 15: 1–27. <https://doi.org/10.22179/REVMACN.15.165>
- Figuerola J (2015) Interacciones humano – oso andino *Tremarctos ornatus* en el Perú: Consumo de cultivos y depredación del ganado. *Therya* 6(1): 251–278. <https://doi.org/10.12933/therya-15-251>
- Goldstein I, Paisley S, Wallace R, Jorgenson JP, Cuesta F, Castellanos A (2006) Andean bear – livestock conflicts: A review. *Ursus* 17(1): 8–15. [https://doi.org/10.2192/1537-6176\(2006\)17\[8:ABCAR\]2.0.CO;2](https://doi.org/10.2192/1537-6176(2006)17[8:ABCAR]2.0.CO;2)
- Gonzales FN, Neira-Llerena J, Llerena G, Zeballos H (2016) Small vertebrates in the spectacled bear's diet (*Tremarctos ornatus* Cuvier, 1825) in the north of Peru. *Revista Peruana de Biología* 23(1): 61–66. <https://doi.org/10.15381/rpb.v23i1.11834>
- Heck V, Figuerola A (2009) Climatología de la zona norte del parque nacional natural Puracé y su análisis en el marco de los datos históricos existentes. In: Figuerola A, Valencia M (Eds) Fragmentación y coberturas vegetales en ecosistemas andinos, departamento del Cauca. Universidad del Cauca, Popayán, Cauca, 57–75.
- Hernández-Guzmán A, Payán E, Monroy-Vilchis O (2011) Hábitos alimentarios del *Puma concolor* (Carnivora: Felidae) en el Parque Nacional Natural Puracé, Colombia. *Revista de Biología Tropical* 59(3): 1285–1294. <https://doi.org/10.15517/rbt.v0i0.3399>
- Hernani-Lineros L, Garcia E, Pacheco LF (2020) Andean bear diet near to and far from a road. *Ursus* (31e7): 1–7. <https://doi.org/10.2192/URSUS-D-19-0003.1>
- Lizcano D, Pizarro V, Cavelier J, Carmona J (2002) Geographic distribution and population size of the Mountain tapir (*Tapirus pinchaque*) in Colombia. *Journal of Biogeography* 29(1): 7–15. <https://doi.org/10.1046/j.1365-2699.2002.00654.x>
- Lizcano D, González TM, Sandoval-A S, Peña J, Galvis C, Mosquera-Guerra F (2017) Características de la danta de montaña (*Tapirus pinchaque*). In: Mantilla-Meluk H, Mosquera-Guerra F, Trujillo F, Lizcano D, Sandoval S, González TM, Londoño MC (Eds) Plan de manejo para la conservación de la danta de montaña (*Tapirus pinchaque*) en el departamento del Quindío. Corporación Autónoma Regional del Quindío (CRQ), Armenia, Quindío, Colombia, 10–15.
- López-Ordóñez JP, Gómez-Lora I, Ramírez-Chaves HE (2020) A record of scavenging behavior of the Andean bear (*Tremarctos ornatus*) on mountain tapir (*Tapirus pinchaque*) in southwestern Colombia. *International Bear News* 29(2): 27–28.
- Márquez R, Goldstein I (2014) Manual Para el Reconocimiento y Evaluación de Eventos de Depredación de Ganado por Carnívoros Silvestres. Wildlife Conservation Society – Colombia. Santiago de Cali, 35 pp.
- Narváez V, Zapata-Ríos G (2016) Guía Para la Identificación de Ataques a Animales Domésticos Causados por Carnívoros Grandes. Ministerio del Ambiente y Wildlife Conservation Society – Ecuador. Quito, 13 pp.
- Ortega-Andrade HM, Prieto-Torres DA, Gómez-Lora I, Lizcano DJ (2015) Ecological and geographical analysis of the distribution of the Mountain Tapir (*Tapirus pinchaque*) in Ecuador: Importance of Protected Areas in future scenarios of Global Warming. *PLoS ONE* 10(3): e0121137. <https://doi.org/10.1371/journal.pone.0121137>

- Padilla M, Dowler RC, Dowler CC (2010) *Tapirus pinchaque* (Perissodactyla: Tapiridae). Mammalian Species 42: 166–182. <https://doi.org/10.1644/863.1>
- Rodríguez A, Gómez R, Moreno A, Cuellar C, Lizcano DJ (2014) Record of a mountain tapir attacked by an Andean bear on a camera trap. Tapir Conservation 23: 25–26.
- Roulin FD (1829) Mémoire pour servir á l'histoire du tapir; et description d'une espèce nouvelle appartenant aux hautes régions de la Cordillère des Andes. Annales des Sciences Naturelle Zoologie Paris 17: 26–55.